

Inside an evaporating two-dimensional charged black hole

Amos Ori

*Department of Physics,
Technion—Israel Institute of Technology, Haifa, 32000, Israel*

We investigate the inner structure of an evaporating charged black hole, within the context of semiclassical dilaton gravity in two dimensions. The matter fields are charged, allowing the evaporation of both the mass and charge of the black hole. We find that the semiclassical effects cause the inner horizon to expand (by a finite factor), rather than to shrink to a point singularity. Although this expansion is a quantum phenomenon, the overall expansion factor is found to be independent of the magnitude of the quantum terms in the effective theory.

The internal structure of charged black holes (BHs) has been the subject of many investigations over several decades. The Reissner-Nordstrom solution admits a perfectly regular inner horizon (IH) inside the BH, as opposed to the spacelike singularity of the uncharged Schwarzschild solution. Classical perturbations are known to develop instabilities on the IH, which lead to a null curvature singularity [1]. However, this singularity is found to be weak and non-destructive, in the sense that its tidal effect on a physical object is bounded and small [2]. A similar situation was observed in a spinning BH [3].

When semiclassical effects are taken into account, the black hole evaporates. In the case of a charged BH there are two distinct semiclassical phenomena: The electromagnetic Schwinger effect, and the gravitational Hawking effect. The evaporation of a charged BH involves both phenomena. An important question is, how does this evaporation affect the internal geometry. The goal of the present paper is to address this problem within a simplified two-dimensional (2d) model.

Callan et al [4] developed a dilatonic 2d model of an uncharged BH. This model as well as similar ones were subsequently investigated by many authors [5]. Similar models of a charged BH (yet with uncharged matter fields) were later developed [6] [7]. Nojiri and Oda [8] then considered a dilatonic model with a charged quantum matter field. In their model, however, the dilaton-electromagnetic coupling took a non-standard form, which led to undesired causal structure of the charged BHs already at the classical level. This problem was later fixed in Ref. [9].

The model [9] consists of gravity in two dimensions, coupled to a dilaton ϕ , electromagnetic field F_{uv} , and N charged matter fields. These fields were taken to be massless chiral fermions, following Nojiri and Oda [8]. The dilaton-electromagnetic coupling was taken as $e^{-2\phi}$ (rather than the coupling $e^{2\phi}$ used in Ref. [8]). The analysis in [9] focused primarily on the evolution of the various fields outside the evaporating BH. In particular the evaporation rate of the BH's mass and charge was determined. In this paper we shall use the same model to explore the evolution of geometry *inside* the BH, with

special emphasis on the dynamics near the IH.

Recently Frolov, Kristjansson, and Thorlacius (FKT) [10] developed a similar model of 2d dilaton gravity coupled to a charged matter field. The main difference is that in Ref. [10] the matter field is bosonized before semiclassical corrections are calculated. Also this field may be either massive or massless. In the massless case the field equations are closely related to those of Ref. [9]. FKT evolved the field equations (for a massless field) numerically in order to explore the dynamics inside the evaporating charged BH. They concluded that the combination of the Schwinger and Hawking effects turns the IH into a spacelike singularity where ϕ diverges to $+\infty$. We employ here the usual translation of the 2d dilatonic terminology to that of a four-dimensional (4d) spherical model, in which the area coordinate is mapped to $e^{-2\phi}$. Then in this language, the inner horizon was found in [10] to shrink to a point singularity.

Previously several authors studied numerically the backreaction of Schwinger's pair-production effect on the internal structure of a charged BH, in either 4d [11] [12] or 2d gravity [13]. These analyses, however, did not incorporate the Hawking effect and hence the BH did not evaporate, a fact which drastically changes the global structure. Also Balbinot and Brady [14] studied the backreaction to the Hawking effect inside a 2d charged BH, but the Schwinger effect was not included, and the BH was a non-evaporating one. To the best of our knowledge Ref. [10] is the only work published so far which analyzes the inner structure of an evaporating semiclassical charged BH.

In this paper we explore the field equations of Ref. [9] analytically, and reveal the internal structure of a semiclassical charged BH which evaporates its both mass and charge. We find a behavior rather different than that reported in [10]. On approaching the IH, the semiclassical effects cause the area coordinate to *expand* (to a finite value), rather than to shrink to a point singularity. This semiclassical expansion is linear in the Eddington coordinates. These analytical results were also confirmed by numerical simulations of our field equations. The origin of the discrepancy between FKT's analysis and our results still needs to be understood. In this paper we briefly

present our analysis and describe the main results. A more detailed description will be given in a separate paper [15].

The classical action and the semiclassical corrections are given in Ref. [9] (it is the same as in Ref. [8], apart from the modified dilaton-electromagnetic coupling). Our notation here follows that of Ref. [9], except that we choose our units such that the cosmological constant $\lambda = 1$. We express the metric in double-null coordinates, $ds^2 = -e^{2\rho} du dv$. (Both u and v are future-directed, as usual, and the lines of constant u or constant v are respectively parallel to past or future null infinity.) The electromagnetic field is expressed as $F_{uv} = (g/\sqrt{6})e^{2(\phi+\rho)}q$, where $g > 0$ denotes the electromagnetic coupling constant. The system thus consists of three unknowns, ϕ, ρ, q . We then transform to new variables $R \equiv e^{-2\phi}$ and $S \equiv 2(\rho - \phi)$. The resulting field equations [16] include a set of three hyperbolic equations for the three unknowns R, S, q :

$$R_{,uv} = (q^2/R^2 - 1)e^S - K\rho_{,uv}, \quad (1)$$

$$S_{,uv} = -2(q^2/R^3)e^S + K\rho_{,uv}/R, \quad (2)$$

$$q_{,uv} = -Kg^2(q/R^2)e^S, \quad (3)$$

where $K \equiv N/24$ (representing the magnitude of the semiclassical effects), and $\rho = (S - \ln R)/2$. In addition there are two constraint equations, which both take the form

$$R_{,ww} - R_{,w}S_{,w} + \hat{T}_{ww} = 0, \quad (4)$$

where hereafter w stands for either u or v , and

$$\hat{T}_{ww} = K[(\rho_{,ww} - \rho_{,w}^2) + (Kg)^{-2}q_{,w}^2 + \hat{z}_w(w)] \quad (5)$$

represents semiclassical fluxes of energy in the u or v directions. The functions $\hat{z}_u(u)$ and $\hat{z}_v(v)$ are determined from the boundary conditions.

The classical equations are obtained by formally setting $K = 0$. Then $\hat{T}_{ww}$ vanishes, and q is set to a constant. The general solution then forms a two-parameter set, parametrized by the mass m and the charge q , to which we shall refer here as the two-dimensional Reissner-Nordstrom (2dRN) solution. This solution is most easily expressed in diagonal coordinates [6]. For the present analysis, however, we shall need the classical solution in the double-null coordinates u, v .

A key function which appears in the classical solution is $H(R) \equiv R - 2m + q^2/R$. We are considering here the non-extreme case, $m > q > 0$. Then this function has two zeros, at $R_{\pm} = m \pm (m^2 - q^2)^{1/2}$. Correspondingly the charged BH admits two horizons: an event horizon (EH) at $R = R_+$ and an IH at $R = R_-$. The latter also functions as a Cauchy horizon (CH) for initial hypersurfaces outside the BH.

For the analysis below we shall need the classical solution in between the two horizons, namely at $R_- < R <$

R_+ , in the "Eddington-like" gauge in which both R and S depend on the single variable $x \equiv u + v$. The solution consists of the two functions $R(x)$ and $S(x)$, which satisfy

$$e^S = -H(R), \quad R_{,x} = H(R). \quad (6)$$

$R(x)$ is formally expressed through its inverse function,

$$x(R) = \int_m^R [1/H(R')] dR'.$$

(The lower integration limit was set here to m for concreteness.) Note that H is negative at $R_- < R < R_+$, and correspondingly x tends to $-\infty$ at the EH and to $+\infty$ at the IH. In a classical collapse scenario the EH (IH) is located at $u \rightarrow -\infty$ ($v \rightarrow +\infty$) and finite v (u).

We also define the two parameters

$$\kappa_{\pm} \equiv \frac{1}{2}(dH/dR)_{R=R_{\pm}} = \pm[(m^2 - q^2)^{1/2}/R_{\pm}].$$

Note that in our notation $\kappa_+ > 0$ and $\kappa_- < 0$, and that $\kappa_{\pm}$ depend solely on q/m .

The domain $R_- < R < R_+$ is naturally divided into three regions: (i) The *near-EH zone*, namely the region of very negative x ($-x \gg 1$), (ii) The *near-IH zone*, $x \gg 1$, and (iii) The *central zone*, namely the region of moderate x values. In both the near-EH and near-IH zones e^S is exponentially suppressed, like $e^{2\kappa_+x}$ and $e^{2\kappa_-x}$ respectively.

Consider next the subset of classical solutions with fixed ratio $q/m = \alpha$. It depends on a single parameter m . One can easily verify that all solutions in this one-parameter family are related by a simple scaling rule:

$$R(x; m) = mR_0(x), \quad S(x; m) = \ln m + S_0(x),$$

where $R_0(x) \equiv R(x; m = 1)$ and $S_0(x) \equiv S(x; m = 1)$ are the functions corresponding to unit mass (and $q = \alpha$).

As long as the BH is macroscopic ($m \gg K$), the semiclassical effects outside the BH are weak in a local sense; Namely, in the neighborhood of each point the metric is that of a classical 2dRN solution weakly perturbed by the semiclassical fluxes. These weak fluxes accumulate over long time scales, however, and lead to slow decay of m and q . We denote the slowly-drifting values of m and q , measured along the EH, by $m_{eh}(v)$ and $q_{eh}(v)$ respectively. In Ref. [9] we calculated the rate of change of these quantities, and analyzed their long-term evolution. It was shown that for $g < 1/2$ the ratio q/m approaches a constant

$$\alpha = \sqrt{1 - 2g}/(1 - g). \quad (7)$$

Furthermore, if q/m is initially α , then this ratio is preserved during the evaporation. In this case the mass and

charge evaporate at constant rates (apart from corrections of order K/m_{eh} [17]). Setting $v = 0$ at the moment when (by extrapolation) m_{eh} vanishes, we may write

$$m_{eh}(v) = -\beta v, \quad q_{eh}(v) = -\alpha\beta v, \quad (8)$$

where $\beta = Kg^2m/R_+$.

In what follows we shall consider the case $g < 1/2$. For simplicity we shall also assume that the BH's initial q/m is the preferred value (7) (but our main conclusions do not crucially depend on this assumption [15]).

We turn now to discuss the evolution inside the BH, assuming that the latter is macroscopic (throughout the rest of this paper by "macroscopic" we mean $R_- \gg K$, which also implies $m \gg K$). In fact the above statement about the slow semiclassical evolution also holds inside the BH at $R > R_-$ (except at the very neighborhood of R_- , as we discuss below). We can then express the solution for R, S, q in a perturbative manner, where the leading order is the classical 2dRN solution with slowly-varying parameters m and $q = \alpha m$. We shall refer to this approximation as the *adiabatic approximation*. In the range $R_- < R < R_+$ which concerns us here, this expansion takes the form

$$\begin{aligned} R(u, v) &= m_{eh}[R_0(u+v) + \varepsilon R_1(u, v) + O(\varepsilon^2)], \\ S(u, v) &= \ln(m_{eh}) + S_0(u+v) + \varepsilon S_1(u, v) + O(\varepsilon^2), \\ q(u, v) &= m_{eh}[\alpha + \varepsilon q_1(u, v) + O(\varepsilon^2)], \end{aligned} \quad (9)$$

where $\varepsilon \equiv K/m_{eh}$ and, recall, $m_{eh} = -\beta v$. The field equations (1-4) are then automatically satisfied at the leading order ε^0 . At the next order ε^1 one obtains a set of linear field equations for the three new functions $F_1(u, v)$, where hereafter F will stand for the three unknowns R, S, q . The initial data for F_1 at the EH are discussed in Ref. [15].

For the present analysis we do not really need the specific solution for the perturbation fields F_1 . All that we need is to make sure that these fields are well controlled inside the BH (and hence the perturbation term $\propto \varepsilon$ is negligible compared to the leading classical term). A close look at the field equations for F_1 reveals [15] that indeed these fields are well controlled in the near-EH and central zones; However, in the near-IH zone (namely at large positive $u+v$) R_1 and q_1 evolve linearly in u and v (see below) and get large values. This linear drift formally invalidates the above perturbative scheme at large positive $u+v$. Since in this paper we are primarily interested in the behavior near the IH, we shall develop now another perturbative scheme which will be valid at large positive $u+v$ despite the linear drift in R and q .

We first bring Eqs. (1-3) to a standard hyperbolic form by isolating $R_{,uv}$ and $S_{,uv}$. We obtain a system of the form

$$F_{,uv} = A_F(R, q)e^S + KB_F(R)R_{,u}R_{,v}, \quad (10)$$

with $A_q = -Kg^2q/R^2$ and $B_q = 0$. For $F = R, S$ the coefficients A_F, B_F are certain functions of their arguments (and K) which we write explicitly in Ref. [15]. For the present analysis we shall not need the B -functions, and also it will be sufficient to express A_R and A_S at the leading order K^0 :

$$A_R \simeq q^2/R^2 - 1, \quad A_S \simeq -2q^2/R^3.$$

In Ref. [15] we analyze the effect of the terms proportional to $R_{,u}R_{,v}$ on the near-IH asymptotic behavior and find it to be negligible (note that $R_{,w}$ vanishes at the IH at the leading order). Omitting these terms, the evolution equations take the simple form

$$F_{,uv} = A_F(R, q)e^S. \quad (11)$$

Our goal is to explore the asymptotic behavior of this system at large $u+v$. To this end we first consider the classical case (namely $K = 0$ and $q = \text{const}$). From Eq. (6) it follows that S decreases at large x linearly, $S \approx 2\kappa_-(u+v)$. Consequently e^S becomes negligible, and Eq. (11) reduces to

$$F_{,uv} \simeq 0. \quad (12)$$

The above asymptotic form of S obviously satisfies this equation. The equation for R implies $R \cong R_u(u) + R_v(v) + \text{const}$, and it is indeed satisfied with $R_u = R_v = 0$ and $\text{const} = R_-$.

Turn now again to the system (11) in the semiclassical case. The most crucial ingredient in the above classical asymptotic behavior is the global decaying factor e^S . There is no reason to expect that this behavior will be sensitive to small modifications in the coefficients A_F . Thus, the system (11) seems to admit a rather generic class of asymptotic solutions at large $u+v$, in which S decreases linearly (or alike) in u and v , leading to Eq. (12). Then the asymptotic solution should take the simple form

$$F \cong F_u(u) + F_v(v) \quad (13)$$

(up to exponentially-small corrections). The claim that our system of evaporating BH indeed falls into this class of asymptotic solutions will be justified below, by determining the actual functions F_u and F_v .

The functions F_u, F_v must satisfy certain consistency conditions in order to enable this simple asymptotic behavior. First, S_u and S_v must decrease sufficiently fast in u and v respectively (a linear decrease, as in the classical solutions, is sufficient). Second, since the functions A_F diverge at small R (in fact A_F has poles at $R = 0$ and $R = K$), the functions R_u and R_v must have a form which guarantees that R will increase (or at least not decrease) with u and v .

In fact, the analysis below indicates that the asymptotic solution (13) takes the more specific form

$$F \simeq a_F u + b_F v + c_F, \quad (14)$$

where a_F, b_F, c_F are certain constants. In terms of these parameters, the above consistency conditions reduce to (i) $a_S, b_S < 0$, and (ii) $a_R, b_R \geq 0$. The sign of a_q and b_q is unimportant. Note that with such a linear behavior, at large $u + v$ the source terms in Eqs. (11) are all bounded by $(u + v)^2 e^{(a_S u + b_S v)}$, which indeed has a negligible effect and justifies the approximation (12,13).

The determination of the required functions $F_w(w)$ is done by matching the near-IH approximation (12) to the adiabatic expansion (9). This matching is done in the *intermediate zone*

$$1 \ll u + v \ll R_-/K. \quad (15)$$

The inequality $u + v \gg 1$ guarantees that $e^S \propto e^{2\kappa_-(u+v)} \ll 1$ and the near-IH approximation (12) is valid. The inequality $u + v \ll R_-/K$ means that in the adiabatic expansion (9) the terms F_1 (which are $\propto K$ but undergo linear drift at large $u + v$) are unimportant, and the adiabatic approximation holds. A consistency condition for this matching is that in the intermediate zone $F_{,u}$ must be independent of v , and vice versa. If in addition $F_{,w}$ is independent of w , then the linear form (14) holds as well.

The parameters a_S and b_S are nonvanishing already at the classical level ($a_S = b_S = 2\kappa_-$). The semiclassical effects lead to $O(\varepsilon)$ corrections to these parameters, but these small corrections are not needed here. On the other hand, a_R, b_R, a_q, b_q all vanish at the classical level, and our goal is to calculate these four parameters at order ε^1 . It will be sufficient to calculate c_F at order ε^0 , because the linear drift in F is not sensitive to these constants.

Consider first the semiclassical evolution of q . Integrating the field equation (11) for $F = q$ with respect to u or v , yields

$$q_{,w} \cong q_{,w}^{(eh)} - Kg^2 \int_{-\infty}^{\infty} (qe^S/R^2) d\hat{w},$$

where $\hat{w}$ is the null coordinate other than w (e.g. $\hat{w} = v$ when w stands for u), and the integration is done along a line of constant w . Since we only need $q_{,w}$ at order K^1 , it is sufficient to evaluate the integrand at the leading order ε^0 , namely it can be taken as $\alpha e^{S_0}/R_0^2$. Taking the upper integration limit as ∞ is justified because of the exponential decay of e^S at large $u + v$. The integration is then straightforward and the integral becomes $q/R_- - q/R_+$. [Note that unlike q and $R_{\pm}$, $q/R_{\pm}$ is independent of m_{eh} and only depends on α .] Noting that $q_{,u}^{(eh)} = 0$ (e.g. from regularity) and $q_{,v}^{(eh)} \equiv q_{eh,v}$ is $-\alpha\beta = -Kg^2q/R_+$, we obtain at the intermediate zone

$$q_{,u} \cong Kg^2(q/R_+ - q/R_-) \equiv a_q, \quad q_{,v} \cong -Kg^2q/R_- \equiv b_q.$$

Both quantities are independent of v and u , confirming the linear form (14) for q .

Next consider the semiclassical evolution of R . Here again we evaluate $R_{,u}$ and $R_{,v}$ at the intermediate zone.

To this end we use the constraint equation (4). Since we only need $R_{,w}$ at order K^1 , it will be sufficient to calculate $\hat{T}_{ww}$ at its leading order—namely that of a test charged field in a classical 2dRN background. This test-field $\hat{T}_{ww}$ is to be evaluated at the near-IH zone of this 2dRN geometry, where $\rho_{,ww} \cong 0$ and $\rho_{,w} \cong \kappa_-$. Also the boundary conditions imply [9] $\hat{z}_u = \kappa_+^2$ and $\hat{z}_v = 0$. One then finds that $\hat{T}_{uu}$ and $\hat{T}_{vv}$ are just constants. Equation (4) then yields $R_{,w} \cong \hat{T}_{ww}/S_{,w}$ (the corresponding homogeneous solution $R_{,w} = e^S$ may be ignored at large $u + v$). Substituting $S_{,w} \cong 2\kappa_-$ we find

$$R_{,u} \cong K[g^2(q/R_- - q/R_+)^2 + (\kappa_+^2 - \kappa_-^2)]/2\kappa_- \equiv a_R$$

and

$$R_{,v} \cong K[(gq/R_-)^2 - \kappa_-^2]/2\kappa_- \equiv b_R.$$

Again, both $R_{,u}$ and $R_{,v}$ are independent of v and u , which justifies the linear form (14).

As was discussed above, our near-IH approximation will only be valid if both a_R and b_R are positive. Noting that Eq. (7) implies $\kappa_+ = g$, one obtains

$$a_R = Kg^2(R_+/R_- + R_-/R_+), \quad b_R = Kg^2(R_+/R_-).$$

Both parameters are positive as desired.

The constants c_R and c_q are calculated from Eq. (14) by setting $R \simeq R_-^{eh}$ and $q \simeq q_{eh}$ at the intermediate zone, where $R_{\pm}^{eh}$ denotes the value $R_{\pm}$ corresponding to $m_{eh}(v)$. One obtains [9] $c_q = c_R = 0$.

The global picture of the region between the two horizons as emerges from our analysis is fairly simple. Since we have set $v = 0$ at the end-point of evaporation, the CH of the evaporating BH is the line $v = 0$. The EH is located at $u = -\infty$, just as in the non-evaporating case. v is negative and increasing along the EH, and u is positive and increasing along the CH (starting from $u \approx 0$). In both the near-EH and the central zones the spacetime is well described by the adiabatic solution (9), in which the mass and charge slowly evolve with v . In particular, R (like q) decreases linearly along the EH. In the near-IH zone all dynamical variables drift linearly in u and v . Along the CH R (like $-q$) grows linearly, $R \simeq a_R u$, starting from $R \simeq 0$ at $u \simeq 0$ and subsequently growing to macroscopic values.

In considering the near-IH behavior of q , one recognizes that both a_q and b_q are negative. In fact the charge flips its sign before the CH is approached, and along the CH $q \simeq a_q u < 0$.

The positivity of a_R and b_R means that the effect of the semiclassical terms near the IH is to expand the latter (namely to increase R), rather than to shrink it to a point singularity $R = 0$. Consider an infalling observer which crosses the EH at some $v = v_0$ in the macroscopic domain (namely $-v_0 \gg 1$) and takes a typical orbit which in the locally-2dRN background would hit the IH at some finite

u . The observer approaches the central zone at $v \approx v_0$ and $u \approx u_0 \equiv -v_0$. Subsequently on heading towards the IH v increases remarkably but u is essentially frozen at $u \approx u_0$. In the first part of the journey, during the adiabatic phase, R steadily decreases from R_+ to R_- (corresponding to the entrance mass $m_{eh}(v_0)$) just as in a 2dRN spacetime. But subsequently, on approaching the very neighborhood of the CH, R will start increase according to

$$R \simeq a_R|v_0| + b_R v \simeq R_-^{eh}(v_0) + b_R(v - v_0).$$

Note that this phase of increasing R lasts an enormously short proper time, due to the huge blue-shift factor near the IH. The final value at the CH is $R \simeq a_R|v_0|$. A simple calculation shows that this value is even greater than $R_+^{eh}(v_0)$, by the factor $R_+/R_- + R_-/R_+ > 1$.

Note that this final R value at the CH is independent of K ; Namely, although the near-IH increase in R is a quantum phenomenon, its magnitude only depends on the macroscopic parameters m_{eh} and q_{eh} , not on the amplitude of the quantum terms in the effective action. This has a simple explanation: The rate of the near-IH semiclassical increase in R is $\propto K$, but at the same time, since $\beta \propto K$, for a given entrance mass $m_{eh}(v_0)$ the evaporation time (from $m_{eh}(v_0)$ to complete evaporation) is $\propto 1/K$, hence their product is independent of K . The same considerations apply to the semiclassical drift in q , and one finds that the final charge at the CH is $2\kappa_-$ times the entrance value $q_{eh}(v_0)$, independent of K .

We also performed numerical simulations of the full nonlinear hyperbolic system (1-3) [using the form (10) with the exact functions A_F, B_F]. We started from initial data corresponding to Eq. (8) along the EH. These simulations confirm the above analysis. In particular, all quantities F were found to evolve linearly in u and v in the near-IH zone, with R increasing and q decreasing. The numerically computed quantities $F_{,u}$ and $F_{,v}$ all agree with the above analytically calculated parameters a_F and b_F to within a few percents. More details of the numerical simulations will be presented in Ref. [15].

The rapid change in R as a function of the observer's proper time in the near-IH zone raises the question of whether the semiclassical theory is valid in this region. Scalars made of gradients, such as $e^{-2\rho}\phi_{,u}\phi_{,v}$, get enormously large (though finite) values on approaching the CH. Yet the effective gravitational coupling constant e^ϕ [4] remains small (except near $u = v = 0$). Nevertheless the present analysis makes one point very clear: When the semiclassical theory comes to its limit of predictability (either at $v = 0$ or at an earlier stage where gradient-based scalars become large), $R_{,v}$ is *positive* rather than negative. This fact may have important implications to the issue of extension of geometry beyond the CH, which

we shall not address here.

Finally we comment on the relevance of our results to the real 4d world. A key ingredient in the above 2d analysis is the negative sign of $\hat{T}_{uu}$ and $\hat{T}_{vv}$ of a *test* field near the IH. In the case of a 4d spherical charged BH it is possible to bring the semiclassical field equations to a form very similar to the above 2d model. This similarity strongly suggests that in the 4d case too, the crucial factor which will determine whether the CH expands or contracts will be the sign of $\hat{T}_{uu}$ and $\hat{T}_{vv}$ near the IH, for a test quantum field in the Reissner-Nordstrom geometry. Unfortunately in the 4d case there is no simple expression for $\hat{T}_{\mu\nu}$. Nevertheless certain methods were developed for calculating $\hat{T}_{\mu\nu}$ in the Reissner-Nordstrom geometry outside the EH [18]. The present analysis emphasizes the importance of extending the calculation of $\hat{T}_{\mu\nu}$ to the interior of the 4d Reissner-Nordstrom BH, particularly to the neighborhood of the IH.

I am grateful to Larus Thorlacius for warm hospitality and fruitful discussions. I also thank Joshua Feinberg and Joseph Avron for helpful comments.

-
- [1] E. Poisson and W. Israel, Phys. Rev. D **41**, 1796 (1990) and references therein.
 - [2] A. Ori, Phys. Rev. Lett. **67**, 789 (1991).
 - [3] A. Ori, Phys. Rev. Lett. **83**, 5423 (1999) and references therein.
 - [4] C. G. Callan, S. B. Giddings, J. A. Harvey and A. Strominger, Phys. Rev. D **45**, R1005 (1992).
 - [5] See e.g. J. G. Russo, L. Susskind and L. Thorlacius, Phys. Rev. D **46**, 3444 (1992).
 - [6] M. D. McGuigan, C. R. Nappi and S. A. Yost, Nucl. Phys. **B375**, 421 (1992).
 - [7] E. Elizalde and S. D. Odintsov, Nucl. Phys. **B399**, 581 (1993).
 - [8] S. Nojiri and I. Oda, Phys. Lett. **B294**, 317 (1992).
 - [9] A. Ori, Phys. Rev. D **63**, 104016 (2001).
 - [10] A. V. Frolov, K. R. Kristjansson, and L. Thorlacius, Phys. Rev. D **73**, 124036 (2006).
 - [11] I. D. Novikov and A. A. Starobinsky, Sov. Phys. JETP **51**, 1 (1980).
 - [12] E. Sorkin and T. Piran, Phys. Rev. D **63**, 084006 (2001).
 - [13] A. V. Frolov, K. R. Kristjansson, and L. Thorlacius, Phys. Rev. D **72**, 021501 (2005).
 - [14] R. Balbinot and P. R. Brady, Classical Quantum Gravity **11**, 1763 (1994).
 - [15] A. Ori, in preparation
 - [16] See Eqs. (55,61-65,71,72) in Ref. [9].
 - [17] These $O(K/m)$ terms lead to logarithmic corrections in Eq. (8). Nevertheless these corrections are negligible in the macroscopic case and will be ignored here.
 - [18] P. R. Anderson, W. A. Hiscock and D. A. Samuel, Phys. Rev. D **51**, 4337 (1995).